# Missouri Botanical Garden

## Peter H. Raven Library

# George Engelmann Botanical Notebooks



*Pagination Note:*
*Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.*

Trifolium from Wisconsin farm May 30 1870

607



after soaking

x4

legume x4

Mostly 2 seeds

x20





pistil x4

4 ovules

upper teeth equal to others with wide sinus,

Stipel connate adnate to base of petiole

CHAS. HABICHT & CO.,

Successors to

**Enno Sander & Co.,**

# PHARMACEUTISTS,

Walnut & Fourth Sts.,
SOUTHERN HOTEL,

Walnut & Second Sts.,
BARNUM'S HOTEL,

ST LOUIS, MO.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MISSOURI BOTANICAL GARDEN

Trifolieæ — portions

May 30 1870

608

×415



2 ovules

pistil ×4

style very thin below, swollen higher up.

×4

petals & stamens connate ½ long tube





lowest both legume, upper with narrow slit but scarcely connate



legume

×4

×20



stipules acute, connate adnate for ½

CHAS. HABICHT & CO.,

Successors to

**Enno Sander & Co.,**

# PHARMACEUTISTS,

Walnut & Fourth Sts.,
SOUTHERN HOTEL,

Walnut & Second Sts.,
BARNUM'S HOTEL,

ST LOUIS, MO.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MISSOURI BOTANICAL GARDEN

Baptisia leucantha · C09

Sedonia July 20 1871

phyllodia on trunk and frequently thin
axis along branches in 1/3 arrangement
leaves on branches in 1/2, flowers
in compound racemes ord. $\frac{7}{18}$?

There is always a primary (?) underlying
leafbud between leaf and branch, on
stem as well as branches

Missouri Botanical Garden
George Engelmann Papers

*Sir:*

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
Missouri Botanical Garden

*Physostegia venenosum* Balfour
in Trans. Roy. Soc. Edinby. 22
Pascolarcos, pg 305. 2 plates.

Hanbury in Seemann's Journal of Botany
1. p. 239. (Aug 1863)

610

San Francisco Mts. 112.22 west.
35° 12 north

Missouri Botanical Garden
George Engelmann Papers

Leitch & Cories
DRUGGISTS
South West Corner Fifth and Market Streets.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
Missouri Botanical Garden

Nov. 1848.

*Eysenhardtia*.

E. Drummondi H. Baumpfl. Lindheimer ... August - fl.
spike dense multifl
Calyx tubulosus, basi obconicus, glandulosus; dentibus triangulibus, subaequalibus, 2 superioribus paulo latioribus, reliquis oribus infimo non longiore; — corolla papilionacea; vexillo cuneato-oblongo, integro s. paulo emarginato alis oblongo-cuneatis, obtusis; — staminibus monadelphis in basi tubi connatis; ovario oblongo, glabro ovulis bi-ovulato, stylo recto apice uncinato, stigmate ... glandulosum supra spectante; legumine oblongo duplo superante, obovato falcato, stylo recto persistente, semine unico subreniformi.

Foliolis 12 jugis foliolis oblongo-lanceolatis, obtusis s. emarginatis, supra glabris subtus reticulatis, glandulosis et adpresse pubescentibus; stipulis subulatis stipellis brevibus subulatis.

Shrub 4-7 feet high. Leaves 1-2 inches long

611

E. Rinconada, Dr Gregg. May - fl.

calyx similar, only teeth shorter quite rounded; petals vexillum more distinctly emarginate and anthers similar; style and ovary hairy, style hooked. leaves a little larger, more distinctly glandulous crenulate, pubescent on both s. Dec - leaflets 12-14 pairs

Shrub, 3 feet high. Leaves 1½-2 inches long

E. spinosa n. sp. Lake Encinillas Dec 1848 (Chihuahua)
spike pauciflora
calyx obconico-campanulatus, vix glandulosus, puberulus; dentibus triangulatis, obtusis, 2 superioribus latioribus reliquis; infimo vix longiore; corolla papilionacea; petalis longitudine subaequalibus, vexillo bilobo, alis oblongis obtusis staminibus subdidelphis in tubum, fissum decimo inferne alteris longioribus connatis; ovario oblongo 4 ovulato, stylo recto pilosa, apice uncinato, stigmate glandulosum supra spectante; legumine ... foliis 6-8 jugis, foliolis ovatis acutis minutim adpresse pilosis, infra glandulosis, stipulatis stipellatisque

A. rough, very much branched shrub; almost spinous for the persistent pedunculos or rhachis; bark dark brown or black; shrub, 2-3 feet high, fl. ~~white~~ white, at last reddish; leaves 4-6 or 7 l. long, leaflets 1-1¼ l. spikes ~~many~~ about one inch long with very stout rhachis begins to flower End of August (at the commencement of the rainy season.)

Eysenhardtia Mountains of Coniguiriachi Dr. Wislizenus 16 Sept 156.

spikis densa multiflora villoso viridi glandulosa triangulatis,
Calyce campanulato; dentibus superioribus latioribus, remotioribus
infimo longiore angustiore; vexillo orbiculato, cordato; staminibus
monadelphis, tubo supra fisso; ovario villoso 2-ovulato,
stylo filiformi, uncinato-recurvo, stigmate acuto.
infra pilosa supra nuda

Foliis 3-4 jugis, foliolis oblanceolatis, obtusis, basi acutis;
stipulis stipellisque minutis
angustatis (rotundatis ?)

(X ultra quae nigro punctulata glandulosa pilosiusculis que,)

Shrub 2-3 feet high; bark whitish; flowers larger than in any of the other species. — spikes white
2-4 inches long leaflets 8-10 lines by 2-2½ l. wide; besides
the black dots on both sides there are innumerable
transparent glands in the substance of the leaf.
It is distinguished from the three others
by the cordate orbicular vexillum, by the gland on
the anther, by the filiform long hooked style,
by the transparently punctate and on both sides dotted
leaves. ——

N.B. The gland on the style of the 3 first species is
not the stigma, but only a smooth gland; the stigma
itself is terminal ?. obtuse.

(X A small tree, Guanaxato — Nothing of the gland on the style is said.)

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

E. Drummii

612



tube closed

vexill



Papilionaceous flower

E. Greggii



vexill



E. spinosa

vexill



alternately longer



a

a

Tube open at a



E. Wislizeni

vexil

ala

carina



Tube open



tube and stamens will be longer in the fully developed flower — this is for a bird

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

dentibus superior[illegible]

~~deciduis~~; Petala 5 – corollam papilionaceam disposita, vexillo oblongo cucullato carinalibus distinctis; Stamina 10 monadelpha s. diadelpha abbreviato-globoso glandulosa. Ovarium 2 – 4 ovulatum; Stylus filiformis rectus apice curvatus s. uncinatus, ~~glandula~~ sub stigmate obtuso ~~lateralibus~~ lateraliter 1-glandulatus. Legumen compressum oblongo-lanceolatum, lunulatum, rostratum, 1- loculare 1- sperm.

Frutices et arbores ~~boreali~~-mexicanae et texanae ~~omnes~~

Amorphae ~~sparsae~~. Folia impari pinnata, multijuga, ~~alterna~~ nigro- impunctata ~~que et legumina~~ glandulosa; foliola stipellata. Racemi terminales pauciflori s. multiflori; flores albidi. calyces et legumina glandulosa

(If you leave out: dentibus inferioribus subaequalibus) (and calyx – s. carinalis) (this will include so far as I understand it, Lysentelis Amorpha, the style of which is unknown to me.)

1. E. amorphoides HBK. arborea; spicis ~~axillaribus~~ terminalibus, calycibus ~~brevi~~ obconico-campanulatis, stam 2. d. 4 Tres typ

2. E. Drummondii Torr. putidosa, foliis 12-jugis, foliolis oblongo-lanceolatis, obtusis s. emarginatis supra glabris, subtus reticulatis glandulosis et adpresse pubentibus, margine ~~ab~~ integris; stipulis subulato setaceis, stipellis brevibus subulatis; spicis paniculato-aggregatis, multifloris, elongatis, calycis dentibus triangulatibus subaequalibus; vexillo ~~oblongo~~ apice integro s. vix emarginato, staminibus tubo brevissimo integro monadelphis; ovario biovulato; ~~et~~ stylo glabro, apice incurvo, ~~glandula~~ ~~[illegible]~~ fl. Jul – Aug.

N. Brazos, Texas, T. Lindheimer, – Shrub 4 – 7 feet high, leaves 1 – 2 inches long; fl. white; style curved upward but not hooked; glands large, ~~Fune~~ directed ~~downwards~~ towards the carina.

3. E. Greggii n. sp. frutescens, foliis 12 – 16 jugis, foliolis oblongis obtusis, utrinque pubescentibus, infra glandulosis margine glandulosa-ciliatis; spicis ~~paniculato aggregatis~~ ut supra, calycibus dentibus abbreviatis rotundatis, subaequalibus, vexillo apice brevi bilobo, ovario biovulato et stylo apice subuncinato pubescentibus.

Rinconada, Northeast Mexico Dr Gregg: fl May; Shrub about 3 feet high; leaves 1½ – 2 inches long; white style almost uncinate, more curved than the filiform spikes; glands directed towards the carina. – Very near E. Drummondii.

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

[illegible]

staminibus, subaequalibus; vexilla profunde bilobo, staminibus
tubo longiore) sub didelphis, ovario 4 ovulato, et stylo
apice uncinato pilosis.

In Lake Enorin Asa north of Okiheehee. fl. August. Sept.
A very remarkable with
in many respects remarkable with
A single looking spumb 2-3 feet high, with black bark
leaflets 1-1¼ l. long; spikes 1 inch long with stout
persistent rhachis; fl. white, turning reddish; upper most
filament, shortest, and almost free, only adhering at base
to the tube; — style strongly hooked, glad large, directed
towards the vexillum (on the hook)

Eysenmannia n gen. calyx campanulate 5 dentate-
dent. sup. Cat. rem. Stamen augustior longior, Pet 5 ÷ or
pap disp; vexilla rotundo orbiculato integro, carin. distinct.
Stamen 10 c. fissum dorsali monadelph v. d. 2 or. — Styles
capillaceus truncato eglandulosus stigmate obliquo-
acuto.
Legume ? —

Frutex mexicanus, ramis Eysenhardtiae fere; folia
imp. pin 3-4 jug. rigro glandulosa et pellucido punctata;
foliola minute stipellata. Race. Terminal, panicled
multiflori; calyces glandulosi; villosi, flores alb.
styles ovar stylosque pilosus.

Distinguished from Eys. by the obovulate ovar,
glandulous anthers, capillary eglandulous style, and
acute stig. but not seen.

E. texicana n sp. Mts S Loriz. & W. Hl. Sept.
Shrubs 2-3 feet high, with whitish bark; leaves
2-4 inches long, leaflets oblanceolate obtuse, 8-10 l. long
2-2½ l. wide; Besides the black glands on pellucid
pellucid dots, Anthers with a large gland at the apex
between the upper cells. — Style flexuous, somewhat twisted
(flexuous) and with a large upper hooked.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

1843

614

stipulae

flores
mag nat


mag. nat.

fructus

mag. nat

Mexicanus

Astragalus perennis, caulibus pluribus, adsurgentibus, ad apicem pilis adpressis pubescentibus; stipulis triangularibus contis petiolo ~~adnascentibus~~ et inter se connatis; foliolis 12 jugis, inferioribus rotundato – superioribus lanceolato – ellipticis, omnibus emarginatis v. obtusis subcarnosis, supra glabernimis, infra pilis simplicibus ~~sub~~pubescentibus; racemis foliis ~~longioribus~~ superantibus, laxis 10–15-floris, floribus lutantibus vel patulis, bracteis pedicello brevioribus, calycibus albo-villosis cylindraceis, supra gibbosis, dentibus ovatis acutis, ~~tertia~~ tubo tertio brevioribus, corolla ochroleuca calycem duplo superante carina apice violacea,

Legumine glabra, carnosa, ovato, acuto
subcompresso, bicarinati, (seminibus
biserialibus)

in pratis siccioribus (Attendorn Cottam) et ad
colles vicinos (Bluffs) e regione St Louis,
ft. Apr. et Mayo.
fruct. Junio.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN



615

Amorpha ovata

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
GEORGE NEUMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Locus: Lamarbidegan 17.

616

Astragalus sect Missouriensis Nutt

Caule suffruticoso stipulis petiolo adnatis ovatis subacutis, junioribus sericeis, 6–10 jugis foliolis parvis, ovatis obtusis emarginatis floribus spicato capitatis longe pedunculatis, pedunc. foliis vix longioribus; floribus violaceis deflexis pedicellatis brevibus glabris (pedicello longitudine); leguminibus glabris arcuatis acuminatis denique erectis, dorso carinato sulcatis.

H. Mai; Missouri, Fort Smith = Little Rock Arkansas Prairies in Clifton Creek Washington.

Astragalus caroliniana, Nutt in Amer Bottom fort Chipol ovat ant caulin, romatis oppositifoliis! legum ovat tumid rostrat

Astragalus missouriensis Nutt! Figure in Amer bottom = Franklin Co Ozark = Mai =

Planta basin versus subherbacea, caule subcrecto 3–10 pollicari, adpresse pubescente, stipulis subulatis distinctis, persistentibus, foliolis 8–12 jugis, ovatis obovatis glabriusculis, supernorum ovato-lanceolatis obtusis puberulis, spicis ovatis pedunculatis, longitud. foliorum & 4–6 floris, floribus laete violaceis vix pedicellatis longitudine; calycis villoso albido pubescentis dentibus ovatis, corolla duplo longioribus; vexillo supra flavo fusco, — legumine inflato globoso.

Astragalus ? alter micranthus Nutt.

caule subcrecto 1–1½ pedali glabro, stipulis sub-caulinis distinctis, ovati acutis, foliis 6–8 jugis, foliolis

4

glongis · emarginatis, glabriusculis, nervis pilosis.
pedunc. fructiferis elongatis duplo folior longitudine,
p. crassiusculis, bracteis lanceol. minutis, floribus 3–6
fasciculatis.

Frutex ℏ-pollen — weiß Blüten in Trauben

Phas. leiospermus T. & Gr. ℘ Nro 13

Phaseolus · helvolus ?!

Nahe verwandtschaft mit dem Ph. vexillatus (vielleicht
dasselbe !!) Stengel oben blauen Filz glatt, ganz
kahl, Blüten gelb, — Schiffchen spitzig — Fahne mit rothen Adern
weiß — Flügel gelb — Fahne mit rothen Adern — Stengel
helvolus (Stengel auf rothen Adern behaart)
(Ph. dasycarpus Stengel rothen Adern behaart)
Blüten weiß Phaseolus leiospermus
nennen.

Rhynchosia

Stengel windend und dünn
Blätter — Fahne erecta pubescens, foliol.
ovatis acutis integris glanduloso
Stengel braun behaart Blüten in Trauben
tomentosa pubescens.

Rh. dasycarpa ! Stengel buschig
Stengel rothen Adern behaart foliol.
infimis integris, superne ternatis rotundo
subrotundis, spicis floralibus foliis longiori
de ramulicia Stengel erecta Nat. Stengel
braun behaart Blüten.



Dalea lanata Spreng Pandjis Arkansas Ozarks; Spreng. Exemplare von Arkansas — Nicht gesehen
by fifth growth and pell pubescens 4–7 jugis
runcato retusis, submucronatis, pellucido=
punctatis utrinque lanatis — Bracteis (non
subulatis!) ovatis acuminatis calycem sub-
aequantibus — — ruolis decumbente dense villosis
flor violaceis. July flowt.
hofs in Arkansas Ozarks!

Viria americana Mchx. Arkansas; Ozarks. 2 glabra foll oblong-oval, obtusa
mucronata, stipul semisagittat. — dentalis fl. viol.
2/ glabra, foliis ovalis acutis mucronatis
stipulis angustis minis semisagittatis
subdentatis flor violaceis.
3/ pubescens foll ovatis obtusis submucro-
natis, stipulis minutis semisagittatis
villosis. — flor b-. purpureis

Viria tridentata ♃ ? Ark. in Ozarks
by Nuttall. Upon Stalk, 3', 4-5
Branches in an Erect, slightly tufted, bifid at integris
racemi pauciflor foliis brevioribus nervosis

Erium ~~pilosum~~ Ozarkanum nov. spec? Found by
Arkansas river; in very dry sandy soil, pilot
in the Ozarks, also dried specimens from
Nuttall in the woods.
Blossoms in July. Stalk 3' high, pilose
above Habitat occasper

Viria microantha Viria microantha T. & G.
Erium microanthum T. & G. nov. spec. Grows
Plains on the Arkansas creek, in Salina, Arkansas

Viria Lemmonii T. & G.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

J. & G.

Yucca mirranthe

in Mai — Am Abg 22 fabelhaft viele Exem.
beb d Post eilt sehr breit wie die Zunge 2 d
dt — 5 Paare j. pedunc 2 Abg hinaus
Post aller Gatt; jedes Auge mit 2–3
jungen Köpfen. Sehr pfl Form, ähnl
pall, Bau 2 — i rasch wild wdg Rüg
fast umgek O — blatt retus bis
obcordat, mucronat

grosse dentalis — superioribus stipul sag Wells
integris. linear sag Hall

Salkeyus praec. Ell. ? stipul eiförm
blattpaarig 1 bis 2 (–11 1=) blattig; stipulae
semiorbiculatae, accretae (fast falcatae)
2 kurz tripl (eiförm-lanz). —
abstehend, hellbraun. Spitze dopp
oft in mehre Fasern
kurz Nutt.

Aloë ? Cooleyi Ell ? Dfst pfl
bis 1 Fuß; mein Pflanz ist sie in Aloë
(od Darlingtonia ?) mit lgpall pfindl Blatt
2 mit kurz pfl blattig — pfindl Dfst
mehr zu gras 2 mit 4 — 6 blattl retus bis
obl mucron in mucron ab eine abgel blattig
Zweig bilden. — Es mein die Pflanz
von Esto ist, ist zu pfl. foliis eglandulosa
bipinnata 10–20 jug — foliolis 25–30 jug
(foliola 2000–2500 !! in folio singulo !!) — will
sie zu geben. Auf Ihre in Buch
Zeit 6–Abr 9–65 — Juni.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Lespedeza stylosanthoides n sp

ovate caudate stipules adnate to stem

base rounded ovary & calyx

pubescent leaves

618



MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN